arXiv:hep-ph/0203224v1 23 Mar 2002

NYU-TH/02-03-01
hep-ph/0203224

Simple Formulae for $\sin^2\theta_{eff}^{lept}$, M_W, Γ_l, and their Physical Applications

A. Ferroglia[a,1], G. Ossola[a,2], M. Passera[b,3], and A. Sirlin[a,4]

[a] *Department of Physics, New York University,*
4 Washington Place, New York, NY 10003, USA

[b] *Dipartimento di Fisica "G. Galilei", Università di Padova and*
INFN, Sezione di Padova, Via Marzolo 8, I-35131, Padova, Italy

Abstract

The calculation of the partial widths of the Z boson in the effective scheme is discussed. Simple formulae for $\sin^2\theta_{eff}^{lept}$, M_W, and Γ_l in the on-shell, $\overline{MS}$, and effective renormalization schemes are presented. Their domain of validity extends to low M_H values probed by current experiments, and to the range of $\Delta\alpha_h^{(5)}$ results from recent calculations. Physical applications of these formulae are illustrated. A comparative analysis of the above mentioned schemes is presented. The paper includes a discussion of the discrepancy of the $\sin^2\theta_{eff}^{lept}$ values derived from the leptonic and hadronic sectors and of the shortcomings of theoretical error estimates of $\sin^2\theta_{eff}^{lept}$ based solely on those affecting M_W.

---

[1] e-mail: andrea.ferroglia@physics.nyu.edu
[2] e-mail: giovanni.ossola@physics.nyu.edu
[3] e-mail: passera@pd.infn.it
[4] e-mail: alberto.sirlin@nyu.edu

1 Introduction

The on-shell [1,2] and $\overline{MS}$ formulations [3,4,5,6,7] are two of the most frequently employed schemes of renormalization in the evaluation of electroweak corrections. The on-shell approach (OS) is very "physical", as it identifies the renormalized parameters with observable, scale-independent constants, such as α, M_W and M_Z. On the other hand, $\overline{MS}$ calculations exhibit very good convergence properties. In fact, they follow closely the structure of the unrenormalized theory and, for this reason, avoid large finite corrections that are frequently induced by renormalization. They also involve inherently scale-dependent parameters, such as $\hat{\alpha}\,(\mu)$ and $\sin^2\hat{\theta}_W\,(\mu)$, which play a crucial role in the analysis of grand-unification.

Recently, a novel approach, the effective scheme of renormalization (EFF), has been proposed [8,9]. It shares the good convergence properties of the $\overline{MS}$ formulation but, at the same time, eliminates the residual scale dependence induced, in finite orders, by the truncation of the perturbative series. Like the OS approach, it identifies the renormalized parameters with observable, scale-independent constants, such as α, $s^2_{eff} \equiv \sin^2\theta^{lept}_{eff}$, and M_Z. Thus, in the OS, $\overline{MS}$, and EFF schemes, the basic electroweak mixing parameter is identified with $\sin^2\theta_W \equiv 1 - M_W^2/M_Z^2$, $\sin^2\hat{\theta}_W\,(\mu)$, and s^2_{eff}, respectively.
It has been shown that the incorporation of the $\mathcal{O}(\alpha^2 M_t^2/M_W^2)$ contributions [10] greatly reduces the scheme and scale dependence of the radiative corrections in the theoretical evaluation of s^2_{eff} and M_W, as well as the upper bound on M_H [11,12]. In these papers the calculations were carried out in two different implementations of the on-shell scheme of renormalization, denoted as OSI and OSII, as well as in the $\overline{MS}$ framework. Comparison of the three calculations led to an analysis of the scheme dependence and, by inference, to an estimate of the theoretical error due to the truncation of the perturbative series. Ref. [13] extended these studies to the analysis of the partial leptonic widths Γ_l of the Z boson. In turn, Refs. [8,9] discussed the results of the s^2_{eff} and M_W calculations in the EFF scheme.

One of the objectives of this paper is to present simple analytic formulae for the precisely measured parameters s^2_{eff}, M_W, and Γ_l in the four schemes ($\overline{MS}$, OSI, OSII, EFF), as functions of M_H and important inputs such as M_t, $\Delta\alpha_h^{(5)}(M_Z)$, and $\alpha_s(M_Z)$, and to illustrate their usefulness in physical applications. Expressions of this type have been discussed before in the framework of the OS and $\overline{MS}$ schemes [12,13]. In the present paper we extend the range of validity of the formulae to lower values of M_H probed by recent measurements, and include the corresponding expressions in the EFF scheme.

The plan of the paper is the following: Section 2 outlines the theoretical derivation of Γ_l in the EFF scheme, which has not been discussed in the literature. Section 3 presents simple formulae for s^2_{eff}, M_W, and Γ_l in the four schemes and shows how to apply them in order to obtain information about M_H and to predict M_W. A discussion of the discrepancy of the s^2_{eff} values derived from the leptonic and hadronic sectors is included. Section 4 presents a comparative analysis of the results in the four schemes and a discussion of the theoretical errors associated with the s^2_{eff} and M_W calculations.

2 Leptonic Partial Widths of the Z boson in the EFF scheme

Theoretical calculations of the partial widths of the Z boson into leptons and the u, d, s, c quark channels that incorporate the corrections of order $\mathcal{O}\left(g^4 (M_t^2/M_W^2)^n\right)$ $(n = 1, 2)$ have been carried out in Ref. [13] in the framework of the $\overline{MS}$, OSI, and OSII schemes. In particular, simple analytic formulae for the leptonic partial widths Γ_l have been presented in the same work.

In this Section we extend the analysis of Γ_l to the EFF scheme. As in Refs. [8,11,13] we retain complete one-loop corrections and two-loop effects enhanced by factors $\left(M_t^2/M_W^2\right)^n$ $(n = 1, 2)$ or involving the product of one-loop imaginary parts. In the latter case, the rationale is that such products are quite sizable, relative to typical $\mathcal{O}(g^4)$ contributions, since imaginary parts involve several additive terms.

It is convenient to start with the $\overline{MS}$ expression [6,13]:

$$\Gamma^l_Z = \frac{\hat{e}^2}{\hat{s}^2 \hat{c}^2} \frac{M_Z}{96\pi} |\hat{\eta}_l| \left\{ 1 - 4|Q_l|\hat{s}^2 Re\ \hat{k}_l + 8 Q_l^2 \hat{s}^4 |\hat{k}_l|^2 \right\} + \mathcal{O}\left(\frac{m_l^2}{M_Z^2}\right) , \quad (1)$$

where $\hat{e}^2$ and $\hat{s}^2 = 1 - \hat{c}^2$ are abbreviations for the $\overline{MS}$ parameters $\hat{e}^2(\mu)$ and $\sin^2 \hat{\theta}_W(\mu)$, respectively, Q_l is the charge of the final lepton, m_l is its mass, and $\hat{k}_l(\mu)$ and $\hat{\eta}_l(\mu)$ stand for appropriate electroweak form factors. QED corrections are included by appending an overall factor of $1 + 3\alpha/4\pi$. An expression analogous to Eq. (1) is valid for the four lightest quarks provided a color factor $N_c = 3$ is appended and $Q_l \hat{k}_l$ is replaced by $Q_f \hat{k}_f$ (f=u-c). In this case, one-loop QCD corrections are included by inserting a factor $1 + \alpha_s(M_z)/4\pi$. These expressions are conventionally evaluated at $\mu = M_Z$ but, in principle, they may be studied over a range of μ-values since the truncation of the perturbative series induces, in general, a residual scale dependence.

As in Ref. [8], we attempt to express Eq. (1) in terms of scale-independent constants such as G_μ, M_Z, and s^2_{eff}. Recalling the basic relation [7]

$$s^2_{eff} = \hat{s}^2 Re\ \hat{k}_l \,, \tag{2}$$

we note that, through terms of $\mathcal{O}(g^4)$, the expression between curly brackets in Eq. (1) becomes

$$1 - 4|Q_l|\hat{s}^2 Re\ \hat{k}_l + 8Q_l^2 \hat{s}^4 |\hat{k}_l|^2 = 1 - 4|Q_l| s^2_{eff} + 8Q_l^2 s^4_{eff} \left[1 + (Im\ \hat{k}_l)^2\right] \,. \tag{3}$$

In the case of the four lightest quarks ($f = u - c$), Eq. (3) is replaced by

$$\begin{aligned} 1 - 4|Q_f|\hat{s}^2 Re\ \hat{k}_f + 8Q_f^2 \hat{s}^4 |\hat{k}_f|^2 &= 1 - 4|Q_f| s^2_{eff} \left[1 + Re(\Delta\hat{k}_f - \Delta\hat{k}_l)\right] \\ &+ 8Q_f^2 s^4_{eff} \left[1 + 2\, Re(\Delta\hat{k}_f - \Delta\hat{k}_l) + (Im\ \hat{k}_f)^2\right] \,. \end{aligned} \tag{4}$$

Next, we turn our attention to the overall coupling $(\hat{e}^2/\hat{s}^2\hat{c}^2)$ and the correction factor $\hat{\eta}_f = 1 + (\hat{e}^2/\hat{s}^2)\Delta\hat{\eta}_f$ in Eq. (1). It is convenient to split the latter into one- and two-loop contributions:

$$(\hat{e}^2/\hat{s}^2)\Delta\hat{\eta}_f = (\hat{e}^2/\hat{s}^2)\Delta\hat{\eta}_f^{(1)} + (\hat{e}^2/\hat{s}^2)^2\Delta\hat{\eta}_f^{(2)} \,. \tag{5}$$

The amplitudes $(\hat{e}^2/\hat{s}^2)\Delta\hat{\eta}_f^{(1)}$ and $(\hat{e}^2/\hat{s}^2)^2\Delta\hat{\eta}_f^{(2)}$ involve the field renormalization of the Z boson and appropriate vertex contributions and have been studied in Ref. [6] and Ref. [13], respectively[5].

In order to express $(\hat{e}^2/\hat{s}^2\hat{c}^2)$ in terms of $G_\mu M_Z^2$, we employ

$$\frac{\hat{e}^2}{\hat{s}^2} = 8\frac{G_\mu}{\sqrt{2}} M_W^2 \left(1 - \frac{\hat{e}^2}{\hat{s}^2}\hat{f}\right) \,, \tag{6}$$

---

[5] In Ref. [6], $1 + (\hat{e}^2/\hat{s}^2)\Delta\hat{\eta}_f^{(1)}$ is denoted as $\overline{\rho_{ff}}$.

which follows from Eq. (20) of Ref. [11]. The amplitude $(\hat{e}^2/\hat{s}^2)\hat{f}$ involves the W-boson self-energy, as well as vertex and box contributions to muon decay, and is discussed in that work. Recalling the relations [1], [5]

$$\frac{M_W^2}{M_Z^2} = \cos^2\theta_W \equiv c^2\,, \tag{7}$$

$$\frac{c^2}{\hat{c}^2} = \frac{1}{1-(\hat{e}^2/\hat{s}^2)\Delta\hat{\rho}}\,, \tag{8}$$

the overall coupling may be expressed in the form:

$$\frac{\hat{e}^2}{\hat{s}^2\hat{c}^2} = 8\frac{G_\mu}{\sqrt{2}}\frac{M_Z^2\left(1-(\hat{e}^2/\hat{s}^2)\hat{f}\right)}{(1-(\hat{e}^2/\hat{s}^2)\Delta\hat{\rho})}\,. \tag{9}$$

In analogy with Eq. (5), it is convenient to split $(\hat{e}^2/\hat{s}^2)\Delta\hat{\rho}$ and $(\hat{e}^2/\hat{s}^2)\hat{f}$ into one and two-loop contributions. Noting that $\Delta\hat{\eta}_f^{(1)}$ and $\hat{f}^{(1)}$ do not contain contributions of $\mathcal{O}\left((M_t^2/M_W^2)^n\right)$ $(n=1,2)$, and expanding in powers of $G_\mu M_Z^2$, we find

$$\begin{aligned}\frac{\hat{e}^2}{\hat{s}^2\hat{c}^2}\,\hat{\eta}_f &= \lambda\Big\{1+\lambda c^2\left[\Delta\hat{\rho}^{(1)}+\Delta\hat{\eta}_f^{(1)}-\hat{f}^{(1)}\right]\\ &+ \lambda^2 c_{eff}^4\Big[\Delta\hat{\rho}^{(2)}+\Delta\hat{\eta}_f^{(2)}-\hat{f}^{(2)}\\ &+ (\Delta\hat{\rho})_{lead}\left(2\Delta\hat{\rho}^{(1)}-(\Delta\hat{\rho})_{lead}+\Delta\hat{\eta}_f^{(1)}-2\hat{f}^{(1)}\right)\Big]\Big\}\,,\end{aligned} \tag{10}$$

where

$$\lambda \equiv \frac{8G_\mu M_Z^2}{\sqrt{2}} \tag{11}$$

and

$$(\Delta\hat{\rho})_{lead} = \frac{3}{64\pi^2}\frac{M_t^2}{M_W^2} \tag{12}$$

is the leading contribution to $\Delta\hat{\rho}^{(1)}$. Since $(\Delta\hat{\rho})_{lead}$ does not involve $\hat{s}^2$, and $\Delta\hat{\eta}_f^{(1)}$ and $\hat{f}^{(1)}$ do not contain terms proportional to M_t^2/M_W^2, we replace $\hat{s}^2 \to s_{eff}^2$ everywhere in Eq. (10). In fact, the difference is a two-loop level effect not enhanced by powers of M_t^2/M_W^2. The functions $\Delta\hat{\rho}^{(i)}$, $\Delta\hat{\eta}_f^{(i)}$, $\hat{f}^{(i)}$ $(i=1,2)$ still depend on both c_{eff}^2 and c^2. Furthermore, the one-loop correction in Eq. (10) is proportional to c^2. As explained in Ref. [8], in order to ensure the complete cancellation of the scale dependence, it is important to express the results in terms of a unique version of the electroweak mixing angle. In order to achieve this objective, we employ the strategy explained in Ref. [8]: i) we replace everywhere M_W^2 by $M_Z^2 c^2$; ii) in the two-loop contributions we substitute $c^2 \to c^2_{eff}$, since the induced difference is of third order; iii) in the one loop expressions we perform a Taylor expansion about $c^2 = c^2_{eff}$, using the one-loop expression for $c^2 - c^2_{eff}$ (cf. Eqs. (9,10) of Ref. [8]). In practice, it is advantageous to evaluate the corresponding derivative numerically, rather than analytically.

Furthermore, since

$$c^2\,(\Delta\hat{\rho})_{lead} = \frac{3}{64\pi^2}\frac{M_t^2}{M_Z^2}$$

is independent of c^2 and the remaining one-loop contributions in Eq. (10), namely

$$\lambda c^2 \left[\Delta\hat{\rho}^{(1)} - (\Delta\hat{\rho})_{lead} + \Delta\hat{\eta}_f^{(1)} - \hat{f}^{(1)}\right] ,$$

are not enhanced by $(M_t^2/M_W^2)^n$ $(n = 1, 2)$, it suffices to replace $c^2 - c^2_{eff}$ by its leading contribution $c^2_{eff} x_t$ $(x_t = 3G_\mu M_t^2/\sqrt{2}\,8\pi^2)$. Substituting Eq. (3) and the result derived from Eq. (10) into Eq. (1), we obtain the final expression for Γ_l in the EFF scheme.

3 Simple formulae for $\sin^2\theta^{lept}_{eff}$, M_W, and Γ_l

In this Section we present simple analytic formulae for s^2_{eff}, M_W, and Γ_l that reproduce accurately the results of the calculations reported in Refs. [8,11,13], and Section 2 of the present paper, over the range $20\,\mathrm{GeV} \leq M_H \leq 300\,\mathrm{GeV}$. We have extended considerably the lower bound of the range since current M_W measurements, by themselves, favor low values of M_H.

The formulae are of the form

$$\begin{aligned} s^2_{eff} &= (s^2_{eff})_0 + c_1 A_1 + c_5 A_1^2 + c_2 A_2 - c_3 A_3 + c_4 A_4, && (13)\\ M_W &= M_W^0 - d_1 A_1 - d_5 A_1^2 - d_2 A_2 + d_3 A_3 - d_4 A_4, && (14)\\ \Gamma_l &= \Gamma_l^0 - g_1\, A_1 - g_5\, A_1^2 - g_2\, A_2 + g_3\, A_3 - g_4\, A_4\,, && (15)\end{aligned}$$

where

$$\begin{aligned} A_1 &\equiv \ln(M_H/100\,\mathrm{GeV})\,, & A_2 &\equiv \left[\Delta\alpha_h^{(5)}/0.02761\right] - 1\,,\\ A_3 &\equiv (M_t/174.3\,\mathrm{GeV})^2 - 1\,, & A_4 &\equiv [\alpha_s(M_Z)/0.118] - 1\,. \qquad (16)\end{aligned}$$

We employ the input parameters [14,15,16]

$$\begin{aligned} M_Z &= 91.1875\,\mathrm{GeV}\,, & \alpha &= 1/137.03599976\,, \\ G_\mu &= 1.16637\times 10^{-5}\,\mathrm{GeV}^{-2}\,, & \Delta\alpha_h^{(5)} &= 0.02761\pm 0.00036\,, \\ M_t &= 174.3\pm 5.1\,\mathrm{GeV}\,, & \alpha_s\left(M_Z\right) &= 0.118\pm 0.002\,, \end{aligned} \tag{17}$$

and include QCD corrections in the $\mu_t = \hat{m}\left(\mu_t\right)$ approach explained in Refs. [11,17]. In contrast with the expressions for s^2_{eff} given in Ref. [12], we note that Eq. (13) contains a term quadratic in A_1, a modification we have introduced in order to extend the accuracy of the formula to low M_H values. Eqs. (13, 14, 15) incorporate complete one-loop corrections, as well as two-loop contributions proportional to $(M_t^2/M_W^2)^n$ $(n = 1, 2)$, studied in Refs. [10,11,13,18], and Section 2 of the present paper.

Tables 1, 2, and 3 present the constants $(s^2_{eff})_0$, M_W^0, Γ_l^0, c_i, d_i, g_i $(i = 1-5)$ in the $\overline{MS}$, OSI, OSII, and EFF schemes. In particular, OSII is strictly independent of the electroweak scale, a property that it shares with the EFF scheme. Over the range $20\,\mathrm{GeV} \leq M_H \leq 300\,\mathrm{GeV}$, and varying the input parameters within 1σ, these formulae reproduce the calculations from the detailed codes with maximum absolute deviations $\approx 10^{-5}$ in s^2_{eff}, $\approx 0.8\,\mathrm{MeV}$ in M_W, and $\approx 7\,\mathrm{KeV}$ in Γ_l.

Although we have employed $\Delta\alpha_h^{(5)} = 0.02761 \pm 0.00036$ [16] in the determination of $(s^2_{eff})_0$, M_W^0, Γ_l^0, c_i, d_i, g_i $(i = 1-5)$, we recall that recent calculations of this important quantity range from 0.027426 ± 0.000190 [19] to 0.027896 ± 0.000395 [20]. For this reason, we have examined the validity of Eqs. (13, 14, 15), with the coefficients given in Tables 1, 2, and 3, over the large range $0.0272 \leq \Delta\alpha_h^{(5)} \leq 0.0283$ that encompasses the recent calculations. We find that the simple formulae still reproduce the results from the detailed codes with maximum absolute deviations that are very close to those reported above.

It is worthwhile to note that the current experimental uncertainty in $G_\mu M_Z^2$ $(\pm 0.0047\%)$ induces tree-level errors of 1.6×10^{-5} in s^2_{eff}, $0.78\,\mathrm{MeV}$ in M_W, and $4\,\mathrm{KeV}$ in Γ_l, which are of the same magnitude as the deviations discussed above. Thus, the accuracy within which Eqs. (13, 14, 15) reproduce the calculations of the detailed codes employed in Refs. [8,11,13] and of the present paper, is sufficient at present.

As a first application, we employ Eq. (13) in the EFF scheme to obtain the value of M_H derived from the current world average $(s^2_{eff})_{exp} = 0.23149 \pm 0.00017$. Noting that $x \equiv c_1 A_1 + c_5 A_1^2$ is normally distributed, we obtain

$x = (1.07 \pm 2.67) \times 10^{-4}$, $x < 5.46 \times 10^{-4}$ at the 95 % confidence level, which leads to

$$M_H = 124^{+82}_{-52}\,\text{GeV}; \quad M_H^{95} = 280\,\text{GeV}, \tag{18}$$

where M_H^{95} stands for the 95 % CL upper bound.

We may also determine M_H using the experimental value of M_W in conjunction with Eq. (14). The current average of the $p\overline{p}$ - collider and LEP2 measurements is $(M_W)_{exp} = 80.451 \pm 0.033\,\text{GeV}$ [14]. Inserting this value in Eq. (14) and calling $y \equiv d_1 A_1 + d_5 A_1^2$, we obtain in the EFF scheme $y = -0.0648 \pm 0.0470$, $y < 0.0117$ at 95 % CL , which corresponds to

$$M_H = 23^{+49}_{-23}\,\text{GeV}\,; \quad M_H^{95} = 122\,\text{GeV}\,. \tag{19}$$

The results given in Eqs. (18,19) have been confirmed on the basis of a χ^2-analysis based on the simple formulae. The central values of M_H have also been verified using the detailed codes.

In summary, for current experimental inputs, M_W leads to significantly lower values of M_H than those derived from s^2_{eff}. This divergence can be also illustrated by using the A_1 value derived from $s^2_{eff} = 0.23149 \pm 0.00017$ and Eq. (13), to predict M_W via Eq. (14). Taking into account the correlation of errors in Eqs. (13, 14), one finds

$$(M_W)_{indir.} = (80.374 \pm 0.025)\ \text{GeV}\,, \tag{20}$$

which differs from $(M_W)_{exp.}$ by $1.86\,\sigma$. It is worth noting that the indirect determination derived by using s^2_{eff} (Eq. (20)) is quite close to the corresponding value $(M_W)_{indir.} = 80.379 \pm 0.023\,\text{GeV}$ obtained in the global analysis [14].

It is also very interesting to obtain M_H using simultaneously the experimental values of s^2_{eff}, M_W, and Γ_l. Employing $(s^2_{eff})_{exp}$ and $(M_W)_{exp}$, and the current world average $(\Gamma_l)_{exp} = 83.984 \pm 0.086\,\text{MeV}$, a χ^2 analysis based on the simple formulae in the EFF scheme leads to

$$M_H = 97^{+66}_{-41}\,\text{GeV}\,; \quad M_H^{95} = 223\,\text{GeV}\,. \tag{21}$$

This result is of course dominated by the s^2_{eff} and M_W data, and may be compared with $M_H^{95} = 196\,\text{GeV}$ in the global fit [14]. The difference is partly due to the fact that $(M_t)_{fit}$ turns out to be somewhat smaller in the global analysis than in our case.

We remind the reader that the current determination of $(s^2_{eff})_{exp}$ shows a χ^2/d.o.f. = 10.6/5, which corresponds to a confidence level of only 6%. Furthermore, the experimental results exhibit an intriguing dichotomy. Those based on the leptonic observables ($A_l(SLD)$, $A_l(P_\tau)$, $A^{(0,l)}_{fb}$) lead to $(s^2_{eff})_l = 0.23113 \pm 0.00021$ while those derived from the hadronic sector ($A^{(0,b)}_{fb}$, $A^{(0,c)}_{fb}$, $< Q_{fb} >$) have an average $(s^2_{eff})_h = 0.23220 \pm 0.00029$. The results within each group agree well with each other but the averages of the two sectors differ by about 3.0σ!

Using Eq. (13) in the EFF scheme, we find that $(s^2_{eff})_l$ leads to

$$M_H = 59^{+50}_{-29}\,\mathrm{GeV}\,; \quad M_H^{95} = 158\,\mathrm{GeV}\,, \tag{22}$$

a result significantly closer to that derived from $(M_W)_{exp}$ (Eq. (19))! The intriguing question remains of whether the difference between $(s^2_{eff})_l$ and $(s^2_{eff})_h$ reflects a statistical fluctuation or is due to new physics involving perhaps the third generation of quarks. In the case of a statistical fluctuation, a possible method of dealing with the discrepancy is to enlarge the error, as discussed in Ref. [12,21]. For instance, if the s^2_{eff} uncertainty is increased by a factor $[\chi^2/\mathrm{d.o.f.}]^{1/2}$ according to the PDG prescription [22], we would have $s^2_{eff} = 0.23149 \pm 0.00025$ which, using Eq. (13) in the EFF scheme leads to

$$M_H = 124^{+104}_{-60}\,\mathrm{GeV}\,; \quad M_H^{95} = 331\,\mathrm{GeV}\,, \tag{23}$$

and

$$M_H = 76^{+64}_{-37}\,\mathrm{GeV}\,; \quad M_H^{95} = 201\,\mathrm{GeV}\,, \tag{24}$$

in the combined s^2_{eff}-M_W-Γ_l analysis. We note that the enlarged error leads to an increase of M_H^{95} from 280 GeV (Eq. (18)) to 331 GeV if we only employed s^2_{eff} as an input, and to a decrease from 223 GeV (Eq. (21)) to 201 GeV in the combined s^2_{eff}-M_W-Γ_l fit. The latter result may be understood as follows: the increasing error in s^2_{eff} changes the shape of the $\chi^2(M_H)$ curve and shifts the position of its minimum to a smaller M_H value. In our case the combination of these two effects leads to a decrease in M_H^{95}. Although this scaling method is not generally employed in current analysis of the electroweak data, it provides a more conservative estimate of s^2_{eff}. It is very interesting to note that it has a small effect in the determination of M_H^{95}, particularly in analyses that combine the s^2_{eff} and M_W data, and that it leads in such a case to a decrease in M_H^{95}! If the discrepancy is due to new physics involving the third generation of quarks [23,24,25] the analysis shows that a substantial, tree-level change in the right-handed $Zb\bar{b}$ coupling is required. Very recently, it has also been argued that, if the $(s^2_{eff})_h$ and $(s^2_{eff})_l$ discrepancy were to settle on the leptonic side, a "new physics" scenario involving light sneutrinos and gauginos would substantially improve the agreement with the electroweak data and the direct lower bound on M_H [26].

4 Comparative analysis of the OS, $\overline{MS}$, and EFF renormalization schemes

As mentioned in the introduction, because of the truncation of the perturbative series, the $\overline{MS}$ calculations have a residual scale dependence, while the EFF scheme is strictly scale independent. This is illustrated in Fig. 1, which compares, in the case $M_H = 100\,\mathrm{GeV}$, the scale dependence of Γ_l in the $\overline{MS}$ scheme with the scale-independent value in the EFF scheme. We note that the latter coincides, to very good approximation, with the maximum of the $\overline{MS}$ curve, which occurs at $\mu \approx 75\,\mathrm{GeV}$, a scale somewhat smaller than M_Z. Nonetheless, the difference between the EFF value and the $\overline{MS}$ calculation at $\mu = M_Z$ is very small.

Tables 4, 5, 6, present the value of s^2_{eff}, M_W, and Γ_l, as functions of M_H, in the four schemes discussed in the paper. They have been obtained using the detailed numerical codes employed in Refs. [8,11,13] and the present paper, the central values of the input parameters in Eq. (17), and $\mu = M_Z$ in the case of the $\overline{MS}$ and OSI schemes. We have kept six significant figures to show more precisely the differences between the various schemes.

It is interesting to note that, in contrast with s^2_{eff} and M_W, Γ_l is not a monotonic function of M_H. Indeed, it shows a maximum at $M_H \approx 52\,\mathrm{GeV}$.

Examination of Table 4 shows that the maximum difference for s^2_{eff} among the four schemes amounts to 3.7×10^{-5} and occurs between OSII and EFF at $M_H = 200, 250, 300\,\mathrm{GeV}$. For M_W, the maximum difference is $2.4\,\mathrm{MeV}$ and occurs between $\overline{MS}$ and OSI at $M_H = 20\,\mathrm{GeV}$. Finally, for Γ_l, the maximum difference is $3.3\,\mathrm{KeV}$ and occurs between OSI and OSII at $M_H = 300\,\mathrm{GeV}$. These differences are quite similar to those encountered in Ref. [8,11] for the s^2_{eff} and M_W calculations over the range $65\,\mathrm{GeV} \leq M_H \leq 600\,\mathrm{GeV}$. They illustrate the scheme dependence of current calculations and provide a first estimate of the theoretical error due to the truncation of the perturbative series. Taking into account further uncertainties associated with the QCD corrections [12,27], the theoretical errors may be enlarged to $\approx 6 \times 10^{-5}$ in s^2_{eff} and $\approx 7\,\mathrm{MeV}$ in M_W.

In the case of M_W, an alternative comparison has been made between OSII, with an M_t-implementation of QCD corrections [11], and important recent calculations that include all $\mathcal{O}(\alpha^2)$ contributions to Δr that contain a fermionic loop [28]. The latter incorporate not only the terms of order $\mathcal{O}(\alpha^2(M_t^2/M_W^2)^n)$ $(n = 1, 2)$, but also a subset of $\mathcal{O}(\alpha^2)$ diagrams not enhanced by powers of M_t^2/M_W^2. Over the range $65\,\mathrm{GeV} \leq M_H \leq 600\,\mathrm{GeV}$, Ref. [28] reported a monotonically decreasing difference between OSII and the more complete calculations, amounting to $\approx 5\,\mathrm{MeV}$ at $M_H = 100\,\mathrm{GeV}$ when the two-loop leptonic contributions to $\Delta\alpha$ are included in both calculations. A glance at Table 5 tells us then that the difference between the other schemes and Ref. [28] is somewhat larger, reaching $\approx 7\,\mathrm{MeV}$ for $\overline{MS}$ at $M_H = 100\,\mathrm{GeV}$.

Since the more complete calculations have not been implemented so far in the case of s^2_{eff} and Γ_l, similar estimates of the theoretical error for these important amplitudes are not yet available. It has been argued [28] that the M_W difference between OSII and the more complete calculations may be used to provide an estimate of the theoretical error in the evaluation of s^2_{eff}, via the OS relation

$$s^2_{eff} = (1 + \Delta k)\, s^2\,, \tag{25}$$

where $s^2 = \sin^2\theta_W = 1 - M_W^2/M_Z^2$ and $1 + \Delta k$ is the relevant electroweak form factor. In this approach, the uncertainty in M_W, and therefore in s^2, is translated into an estimate of the error in the calculation of s^2_{eff}. In our view, such arguments may well be misleading since important corrections involving

$$X = \frac{c^2}{s^2}\, Re\left[\frac{A_{WW}(M_W^2)}{M_W^2} - \frac{A_{ZZ}(M_Z^2)}{M_Z^2}\right] \tag{26}$$

where A_{WW} and A_{ZZ} are the W and Z self-energies, contribute with opposite signs to Δk and Δr [1]. As a consequence, an additional contribution δX to X leads to $\delta s^2 = -[s^2 c^2/(c^2 - s^2)]\delta X$, but only $\delta s^2_{eff} = -[s^4/(c^2 - s^2)]\delta X$! Thus, Δk induces a suppression factor s^2/c^2 in this important class of contributions to s^2_{eff}. The fact that the theoretical uncertainty in s^2_{eff} cannot be simply related to the error in the evaluation of M_W may also be understood by recalling salient features of the $\overline{MS}$ calculation of s^2_{eff}. In the $\overline{MS}$ context, $(1+\Delta k)\, s^2$ in Eq. (25) is replaced by $(1+\Delta\hat{k})\, \hat{s}^2$, an expression that does not explicitly involve M_W at the three level! This feature is even more striking in the EFF calculation of s^2_{eff}, where M_W completely decouples [8]. It is also worth noting that, in the $\overline{MS}$ calculation of s^2_{eff}, the dependence on X does not involve the enhancement factor c^2/s^2. One readily finds that an additional correction δX contributes $\approx -[\hat{s}^4/(\hat{c}^2 - \hat{s}^2)]\,\delta X$ to $(1+\Delta\hat{k})\hat{s}^2$, a result analogous to that found in the OS framework!

5 Conclusions

In Section 2, we have discussed the calculation of the leptonic partial width Γ_l of the Z boson in the effective scheme of renormalization. The extension to the case of the four lightest quarks involves simple modifications, outlined in the text. Our starting point has been the corresponding expressions in the $\overline{MS}$ approach [13]. Both formulations include the corrections of $\mathcal{O}\left(g^4(M_t^2/M_W^2)^n\right)$ $(n = 1, 2)$. A salient feature of the calculations in the effective scheme is that, in contrast with their $\overline{MS}$ counterparts, they are strictly scale independent in finite orders of perturbation theory. This is illustrated in Fig. 1 which depicts, for $M_H = 100\,\mathrm{GeV}$, the scale dependence of current calculations of Γ_l in the $\overline{MS}$ approach and its comparison with the effective scheme result.

In Section 3, we have presented very simple formulae for $\sin^2\theta^{lept}_{eff}$, M_W, and Γ_l, as functions of M_H and important input parameters such as $\Delta\alpha_h^{(5)}$, M_t, and $\alpha_s\,(M_Z)$. A novel feature of these formulae is that their range of validity has been extended to low M_H values probed by current experiments. Furthermore, they retain their accuracy over the range of $\Delta\alpha_h^{(5)}$ values spanned by recent calculations of this important amplitude.

We have illustrated the usefulness of these formulae by extracting M_H and its 95% CL upper bound M_H^{95} employing separately, as inputs, the current experimental averages $(s^2_{eff})_{exp}$ and $(M_W)_{exp}$, as well as the combination of $(s^2_{eff})_{exp}$, $(M_W)_{exp}$, and $(\Gamma_l)_{exp}$. We have also employed the formulae to predict M_W using $(s^2_{eff})_{exp}$ as an input. The result of this prediction turns out to be quite close to that currently derived using the global analysis of all the available electroweak data.

We have also discussed the well-known discrepancy between $(s^2_{eff})_l$ and $(s^2_{eff})_h$, and some of its possible implications for new physics. On the other hand, assuming that the difference is due to statistical fluctuations, we have also considered the effect of enlarging the error according to the well-known

PDG prescription. Surprisingly, we have found that such a procedure decreases M_H^{95} by a small amount in analyses that combine s^2_{eff} and M_W as inputs.

In Section 4, we have compared the calculations of $\sin^2\theta^{lept}_{eff}$, M_W, and Γ_l in the EFF, $\overline{MS}$, OSI, and OSII schemes, over the range $20\,\mathrm{GeV} \leq M_H \leq 300\,\mathrm{GeV}$. We have found maximal differences of 3.7×10^{-5} in $\sin^2\theta^{lept}_{eff}$, 2.4 MeV in M_W and 3.3 KeV in Γ_l. Taking into account uncertainties in the QCD corrections [12,27], the theoretical errors may be enlarged to $\approx 6\times10^{-5}$ in s^2_{eff} and $\approx 7\,\mathrm{MeV}$ in M_W. In the case of M_W, an alternative comparison can be made with important recent calculations that include all $\mathcal{O}(\alpha^2)$ contributions to Δr that contain a fermionic loop [28]. This reveals differences $\approx 7\,\mathrm{MeV}$ at $M_H = 100\,\mathrm{GeV}$ with the four schemes discussed in this paper.

Finally, we have discussed the theoretical shortcomings of estimates of the theoretical error in $\sin^2\theta^{lept}_{eff}$ derived solely from the uncertainty of M_W in the OS scheme.

Acknowledgments

The authors are greatly indebted to G. Degrassi and P. Gambino for very valuable communications. This research was supported in part by the NSF grant PHY-0070787. M.P. acknowledges partial support from the European Program HPRN-CT-2000-00149.

References

[1] A. Sirlin, Phys. Rev. **D22** (1980) 971 and Phys. Rev. **D29** (1984) 89; W. J. Marciano and A. Sirlin, Phys. Rev. **D22** (1980) 2695.

[2] K. Aoki et al., Prog. Theor. Phys. Suppl. **73** (1982) 1; M. Böhm, H. Spiesberger, and W. Hollik, Fortsh. Phys. **34** (1986) 687; W. Hollik, *ibid.* **38** (1990) 165; E. Kraus, Annals Phys. **262** (1998) 155 [arXiv:hep-th/9709154]; P. A. Grassi, Nucl. Phys. **B560** (1999) 499 [arXiv:hep-th/9908188].

[3] A. Sirlin, Phys. Lett. **B232** (1989) 123.

[4] S. Fanchiotti and A. Sirlin, Phys. Rev. **D41** (1990) 319.

[5] G. Degrassi, S. Fanchiotti, and A. Sirlin, Nucl. Phys. **B351** (1991) 49.

[6] G. Degrassi and A. Sirlin, Nucl. Phys. **B352** (1991) 342.

[7] P. Gambino and A. Sirlin, Phys. Rev. D **49** (1994) 1160 [arXiv:hep-ph/9309326].

[8] A. Ferroglia, G. Ossola, and A. Sirlin, Phys. Lett. **B507** (2001) 147 [arXiv:hep-ph/0103001].

[9] A. Ferroglia, G. Ossola, and A. Sirlin, arXiv:hep-ph/0106094.

[10] G. Degrassi, P. Gambino, and A. Vicini, Phys. Lett. **B383** (1996) 219 [arXiv:hep-ph/9603374].

[11] G. Degrassi, P. Gambino, and A. Sirlin, Phys. Lett. **B394** (1997) 188 [arXiv:hep-ph/9611363].

[12] G. Degrassi, P. Gambino, M. Passera and A. Sirlin, Phys. Lett. **B418** (1998) 209 [arXiv:hep-ph/9708311].

[13] G. Degrassi and P. Gambino, Nucl. Phys. **B567** (2000) 3 [arXiv:hep-ph/9905472].

[14] The LEP Electroweak Working Group, EP Preprint Winter 2002 – in preparation (`http://lepewwg.web.cern.ch/LEPEWWG/`).

[15] A. Ferroglia, G. Ossola and A. Sirlin, Nucl. Phys. B **560** (1999) 23 [arXiv:hep-ph/9905442].

[16] H. Burkhardt and B. Pietrzyk, LAPP-EXP-2001-03.

[17] S. Fanchiotti, B. Kniehl, and A. Sirlin, Phys. Rev. **D48** (1993) 307 [arXiv:hep-ph/9212285].

[18] R. Barbieri, M. Beccaria, P. Ciafaloni, G. Curci and A. Vicere, Phys. Lett. B **288** (1992) 95 [arXiv:hep-ph/9205238] and Nucl. Phys. B **409** (1993) 105; J. Fleischer, O. V. Tarasov and F. Jegerlehner, Phys. Lett. B **319** (1993) 249; G. Degrassi, S. Fanchiotti and P. Gambino, Int. J. Mod. Phys. A **10** (1995) 1377 [arXiv:hep-ph/9403250].

[19] A. D. Martin, J. Outhwaite, and M. G. Ryskin, Eur. Phys. J. C **19** (2001) 681 [arXiv:hep-ph/0012231].

[20] F. Jegerlehner, arXiv:hep-ph/0104304.

[21] A. Gurtu, Phys. Lett. **B385** (1996) 415.

[22] Particle Data Group, Review of Particle Physics, Phys. Rev. **D54** (1996) 1.

[23] W. J. Marciano, arXiv:hep-ph/9902332 and Phys. Rev. **D60** (1999) 093006 [arXiv:hep-ph/9903451].

[24] M. S. Chanowitz, arXiv:hep-ph/9905478.

[25] A. Sirlin, in *Proc. of the 19th Intl. Symp. on Photon and Lepton Interactions at High Energy LP99* ed. J.A. Jaros and M.E. Peskin, Int. J. Mod. Phys. **A15S1** (2000) 398 [eConf **C990809** (2000) 398] [arXiv:hep-ph/9912227].

[26] G. Altarelli, F. Caravaglios, G. F. Giudice, P. Gambino, and G. Ridolfi, JHEP **0106** (2001) 018 [arXiv:hep-ph/0106029].

[27] K. Philippides and A. Sirlin, Nucl. Phys. **B450** (1995) 3 [arXiv:hep-ph/9503434].

[28] A. Freitas, W. Hollik, W. Walter, and G. Weiglein, arXiv:hep-ph/0202131.



Figure 1: Scale dependence of Γ_l in the $\overline{MS}$ (dashed line) and EFF (solid line) schemes for $M_H = 100\,\mathrm{GeV}$.

Table 1: Values of the constants appearing in Eq. (13).

Scheme	$\left(s^2_{eff}\right)_0$	$10^4\,c_1$	$10^3\,c_2$	$10^3\,c_3$	$10^4\,c_4$	$10^5\,c_5$
$\overline{MS}$	0.231392	4.908	9.69	2.77	4.5	3.43
OSI	0.231405	4.877	9.69	2.76	4.5	3.34
$OSII$	0.231417	4.999	9.69	2.69	4.4	3.37
EFF	0.231383	4.948	9.69	2.78	4.5	3.50

Table 2: Values of the constants appearing in Eq. (14), in GeV.

Scheme	M_W^0	$10^2\, d_1$	$10\, d_2$	$10\, d_3$	$10^2\, d_4$	$10^3\, d_5$
$\overline{MS}$	80.3868	5.719	5.07	5.42	8.5	8.98
OSI	80.3849	5.667	5.08	5.40	8.5	8.85
$OSII$	80.3847	5.738	5.08	5.37	8.5	8.92
EFF	80.3862	5.730	5.08	5.42	8.5	8.98

Table 3: Values of the constants appearing in Eq. (15), in MeV.

Scheme	Γ_l^0	$10^2\, g_1$	$10\, g_2$	$10\, g_3$	$10\, g_4$	$10^2\, g_5$
$\overline{MS}$	84.0164	4.439	4.77	8.03	1.13	3.670
OSI	84.0174	4.472	4.76	8.04	1.13	3.673
$OSII$	84.0152	4.568	4.76	8.01	1.12	3.683
EFF	84.0176	4.549	4.77	8.05	1.13	3.696

Table 4: Values for s^2_{eff} in the four renormalization schemes considered in the paper. They have been calculated using the central values of the input parameters in Eq. (17).

M_H (GeV)	EFF	$\overline{MS}$	$OSII$	OSI
20	0.230686	0.230699	0.230708	0.230715
50	0.231051	0.231062	0.231080	0.231077
100	0.231385	0.231393	0.231418	0.231406
150	0.231593	0.231599	0.231628	0.231611
200	0.231744	0.231749	0.231781	0.231760
250	0.231864	0.231868	0.231901	0.231878
300	0.231964	0.231967	0.232001	0.231976

Table 5: Values for M_W (in GeV) in the four renormalization schemes considered in the paper (calculated as in Table 4).

M_H (GeV)	EFF	$\overline{MS}$	$OSII$	OSI
20	80.4550	80.4555	80.4539	80.4531
50	80.4219	80.4224	80.4206	80.4202
100	80.3863	80.3869	80.3849	80.3850
150	80.3614	80.3621	80.3600	80.3604
200	80.3421	80.3428	80.3406	80.3413
250	80.3263	80.3270	80.3248	80.3256
300	80.3129	80.3135	80.3114	80.3124

Table 6: Values for Γ_l (in MeV) in the four renormalization schemes considered in the paper (calculated as in Table 4).

M_H (GeV)	EFF	$\overline{MS}$	$OSII$	OSI
20	83.9896	83.9873	83.9878	83.9887
50	84.0353	84.0335	84.0330	84.0346
100	84.0167	84.0155	84.0143	84.0165
150	83.9907	83.9900	83.9882	83.9908
200	83.9667	83.9664	83.9643	83.9672
250	83.9457	83.9457	83.9432	83.9463
300	83.9271	83.9274	83.9247	83.9280